

Hormonal contraception: recent advances and controversies

The Practice Committee of the American Society for Reproductive Medicine

American Society for Reproductive Medicine, Birmingham, Alabama

This document will outline new delivery systems and contraceptive formulations, summarize recent advances in emergency contraception, and review the effects of hormonal contraception on cancer risks, cardiovascular disease, and bone. (Fertil Steril® 2004;82(Suppl 1):26–32. ©2004 by American Society for Reproductive Medicine.)

The first hormonal contraceptive, Enovid™ (150 µg mestranol and 9.85 mg norethynodrel), was approved by the Food and Drug Administration (FDA) for use in the United States in 1960. Oral contraceptives (OCs) are now the most widely used reversible form of hormonal contraception in the United States. A wide variety of hormonal contraceptives are now available. Their mechanisms of action include inhibition of ovulation and/or modification of the endometrium, thus preventing implantation.

In addition to the contraceptive benefits, many other health benefits have been realized with hormonal contraception, including prevention of endometrial and ovarian cancers, control of menstrual bleeding, and relief from cyclic pelvic pain. In the evolution of OCs to their current forms, modifications have been made in efforts to decrease side effects and improve effectiveness and compliance. The first change was a decrease in the dose of estrogen and progestin, which led to the low dose formulations used today (1). Subsequently, new progestins were developed to decrease androgenic side effects. More recently, alternative delivery systems have been introduced in an effort to improve tolerability, compliance, and convenience; these delivery systems include transdermal, implantable, and injectable systems.

This document will outline new delivery systems and contraceptive formulations, summarize recent advances in emergency contraception, and review the effects of hormonal

contraception on cancer risks, cardiovascular disease, and bone.

INJECTABLE COMBINED ESTROGEN/PROGESTIN CONTRACEPTION

The injectable combined estrogen/progestin contraceptive (Lunelle®) was approved by the FDA for use in the United States in 2000. This injectable contraceptive contains 25 mg of medroxyprogesterone acetate and 5 mg of estradiol cypionate (MPA/E₂C) and is administered by deep intramuscular injection every 23 to 33 days. The contraceptive efficacy is similar to depot medroxyprogesterone acetate (DMPA) and OCs, with an estimated failure rate of 0–0.1 per 100 woman-years (2, 3). The most common side effect reported is irregular bleeding during the first three months. Over the longer term, however, most women (83%) have regular menses when using this contraceptive (3, 4). Unlike the other injectable contraceptive DMPA, return to normal ovulatory cycles is rapid and occurs within 63 to 112 days after the final injection (5). MPA/E₂C therefore offers the advantage of an injectable form of contraception with less bleeding and faster return to fertility than DMPA and with a side-effect profile similar to OCs. The product has since been removed from the market and, at the time of this printing, its future was uncertain.

TRANSDERMAL HORMONAL CONTRACEPTION

The transdermal combined estrogen/progestin contraceptive patch (Ortho Evra™/Evra™)

Educational Bulletin

Received March 10, 2004;
 revised and accepted
 March 10, 2004.

No reprints will be
 available.

Correspondence to
 Practice Committee,
 American Society for
 Reproductive Medicine,
 1209 Montgomery
 Highway, Birmingham,
 Alabama 35216.

0015-0282/04/\$30.00
 doi:10.1016/j.fertnstert.2004.
 07.935

was approved by the FDA for use in the United States in 2002. The patch is 20 cm² (4 × 5 cm) and delivers 20 µg/d of ethinyl estradiol and 150 µg/d of norelgestromin (a biologically active metabolite of norgestimate) (6). The dosing is one patch weekly for three consecutive weeks, followed by a patch-free week. The patch may be applied to the buttock, abdomen, upper outer arm, or upper torso excluding the breasts. Mean serum concentrations of hormone are not affected by heat, humidity, exercise, or cold-water immersion (6). The observed contraceptive failure was 0.7 per 100 woman-years (95% CI, 0.31–1.10) but was higher in women with body weight >90 kg (7). The transdermal contraceptive is well tolerated and has a side effect profile similar to OCs (8).

CONTRACEPTIVE VAGINAL RING

A combined estrogen/progestin contraceptive vaginal ring (CVR) (NuvaRing®) was approved by the FDA for use in the United States in 2001. This contraceptive device consists of a flexible ring made of ethylene vinyl acetate copolymer with an outer ring diameter of 54 mm and a cross-sectional diameter of 4 mm. The vaginal ring releases approximately 120 µg of etonogestrel (a biologically active metabolite of desogestrel) and 15 µg of ethinyl estradiol per day (9). The CVR is used for three weeks continuously followed by a one-week ring-free period to allow for regular menstrual bleeding. The advantage for the user is that the CVR is not fitted. It is inserted and removed by the user and may or may not be removed for coitus. The most common side effects reported include headache (6.6%), leukorrhea (5.3%), and vaginitis (5.0%) (10). The failure rate in one study was 0.65 per 100 woman-years (95% CI, 0.24–1.41) (10).

LEVONORGESTREL-RELEASING INTRAUTERINE DEVICE

A levonorgestrel (LNG)-releasing intrauterine device (IUD) (Mirena®) was approved by the FDA for use in the United States in 2000. The first progesterone-releasing IUD (Progestasert®) was approved by the FDA in 1976 but was removed voluntarily from the market in 2001. The LNG-IUD is T-shaped with a steroid reservoir containing 52 mg of levonorgestrel mixed with polydimethylsiloxane, which controls the release rate of hormone. The LNG-IUD remains in place for five years. The LNG-IUD has a failure rate between 0 and 0.2 per 100 woman-years (11); the ectopic pregnancy rate is 0.02 per 100 woman years (12). Menstrual bleeding is decreased by 75% in LNG-IUD users and is attributed to the progestin-induced decidualization and suppression of the endometrium (13); 20% to 50% of users become amenorrheic within the first two years after insertion (14). Following removal, there is rapid return to normal fecundability with one-year life-table pregnancy rates of 89 per 100 for women less than 30 years of age (15).

FERTILITY & STERILITY®

IMPLANTABLE CONTRACEPTIVES

Norplant®, the first implantable contraceptive device, was approved by the FDA for use in the United States in 1990 and withdrawn from the market in 2002 due to complications associated with removal. This device contained 216 mg of levonorgestrel in six implantable rods to be removed after five years. Newer developments in implantable contraception are focusing on fewer implant rods and less androgenic progestins. Norplant® II (Jadelle®) has been approved by the FDA but is not yet marketed in the United States. This system contains two rods 4 cm in length with total dose of 150 mg of levonorgestrel for a three-year duration. The failure rate is 0.8 to 1.0 per 100 woman-years (16, 17). A newer single implant system not yet available (Implanon®) contains the progestin etonogestrel (a desogestrel metabolite having less androgenic activity than levonorgestrel). The carrier polymer, ethylene vinyl acetate, is more stable than the silastic material used in Norplant®.

Among newer implant delivery systems currently under development, biodegradable progestin implants in the form of rods and pellets have the advantage that the implant does not need to be removed (18).

EVOLUTION OF ORAL CONTRACEPTIVES

Oral contraceptive agents have been modified over time to limit estrogen and progestin dosage and decrease side effects. It has been established that progestin combinations with low dose estrogens (30–35 µg ethinyl estradiol) are effective with limited side effects. Lower dose estrogens (20 µg ethinyl estradiol) and newer progestins have subsequently been introduced.

Newer Progestins

The original progestins used in hormonal contraceptives were all derived from ethisterone, an orally active testosterone derivative. Removal of the carbon at C-19 position confers progestational activity, with some residual androgenic activity (19). These progestins are referred to as 19-nortestosterones and include norethindrone, norethynodrel, norethindrone acetate, ethynodiol diacetate, levonorgestrel, and norethindrone enanthate.

The newer progestins include norgestimate, desogestrel, and gestodene (not available in the United States). These progestins were designed to minimize androgenic side effects such as acne, hirsutism, nausea, and lipid changes while increasing progestational effects (19). Among the progestins available in the United States, norgestimate has the greatest progestational effect, and levonorgestrel is the most androgenic (19). The newest progestin, drospirenone, is an analogue of the aldosterone antagonist spironolactone and exhibits both progestational and antiandrogenic activity (20).

Oral Contraceptives Containing Less Than 30 μ g Ethinyl Estradiol

Low dose OCs with 20 μ g ethinyl estradiol were initially marketed for use in perimenopausal women. However, these low dose preparations have been shown to be effective contraceptive agents in women of reproductive age, with reported pregnancy rates ranging between 0.07–2.1 pregnancies per 100 woman-years of treatment (21). When compared with a 35 μ g OC, the 20 μ g OC has comparable cycle control and reduced symptoms of bloating and breast tenderness (22).

Extended OC Therapy

Monthly withdrawal bleeding for women using hormonal contraception has been the traditional prescribing method. Recently, however, this has been challenged and the use of extended hormonal use to delay or prevent menses has been evaluated. A randomized controlled study compared a traditional 28-day cycle to an extended 49-day cycle of a 30 μ g ethinyl estradiol/300 μ g norgestrel monophasic birth control regimen (23). The extended regimen resulted in fewer bleeding days, and no increase in mean spotting or abnormal bleeding episodes. There were no significant differences in other reported side effects such as headaches, weight gain, cramping, or breast tenderness. Similar results have been obtained with extended use of OCs containing 20 μ g of ethinyl estradiol (24).

EMERGENCY CONTRACEPTION

Available hormonal emergency contraceptive (EC) regimens are effective when used within 72 hours of unprotected intercourse regardless of stage in menstrual cycle (25, 26). Either a combined estrogen/progestin regimen or a progestin-only treatment regimen may be used. Both regimens include two doses, the second administered 12 hours after the first. Their effectiveness appears to result primarily from an inhibition or delay of ovulation and neither appears to interrupt or disrupt an already established pregnancy. The recommended combined estrogen/progestin treatment regimen includes 100 μ g of ethinyl estradiol and 1 mg of norgestrel or 0.5 mg of levonorgestrel. The progestin-only regimen involves a higher 0.75 mg dose of levonorgestrel.

There are two FDA-approved and dedicated products for emergency contraception: PrevenTM (pill containing 50 μ g of ethinyl estradiol and 0.25 mg of levonorgestrel; two pills taken 12 hours apart) and Plan B[®] (pill containing 0.75 mg of levonorgestrel; one pill taken 12 hours apart).

The effectiveness of the EC is determined by comparing the number of pregnancies observed following treatment to the expected number of pregnancies in the absence of treatment. In evaluating the combined EC, a meta-analysis involving over 3,000 patients observed a 74% reduction in the pregnancy rate, compared to the theoretical or expected pregnancy rate (27). Less information is available regarding

the success of the progestin-only regimen. The World Health Organization conducted a randomized trial in 1,001 women comparing the progestin-only regimen to the combined regimen (28). The proportion of pregnancies prevented in the progestin-only and combined regimens was 85% and 57%, respectively, with the progestin-only regimen being significantly more effective (RR = 0.36; 95% CI, 0.18–0.70). However, it must be noted that the emergency contraceptive is still less effective in pregnancy prevention than consistent use of other contraceptive methods. Nausea and vomiting are the predominant side effects occurring in 42% and 16% of patients using the combined regimen, respectively (29), though these symptoms are significantly less with the progestin only regimen (28). Ectopic pregnancy can occur following emergency contraception, but the risk does not appear to be increased (30). There are no contraindications to the EC except pregnancy, although no studies have investigated outcomes in women with contraindications to combined OC (31). The progestin-only regimen may be a better choice for women with a personal or a family history of thrombosis.

Antiprogesterins have been evaluated for use as emergency contraception. Mifepristone in a single dose of 600 mg, 100 mg, 50 mg, or 10 mg is equally effective in the prevention of pregnancy and has been shown to be more effective than the combined OC regimen with fewer side effects (32–34).

HORMONAL CONTRACEPTION AND CANCER RISK

Breast Cancer

Breast cancer is very rare in young women. The cumulative risk is less than 10 per 10,000 in women of all races up to age 35 years (35). Breast cancer risk associated with OC use has been evaluated in a pooled analysis from 54 studies involving 53,297 women with breast cancer and over 100,000 controls (36). The main finding was a small increase in the relative risk of localized breast cancer associated with current OC use (RR = 1.24; 95% CI, 1.15–1.33) and also with OC use within one to four years (RR = 1.16; 95% CI, 1.08–1.23), compared to controls. This risk declines shortly after stopping use and disappears within 10 years. By age 50, there is no difference in risk of breast cancer in ever users of OC and controls.

The study also demonstrated that breast cancers diagnosed in OC users were significantly less advanced than those in never users (for spread of disease beyond the breast, RR = 0.88; 95% CI, 0.81–0.95). Overall, there is no evidence of any increase in lifetime risk of breast cancer among OC users. Breast cancer risk associated with use of OC in perimenopausal women has not been evaluated.

Endometrial Cancer

Oral contraceptive use is associated with a lower risk of endometrial cancer. A meta-analysis found that the incidence

would be reduced by 56%, 67%, and 72% with use of combined OCs for four, eight, and 12 years ($P < 0.0001$) (37). The lowered risk of disease persists after stopping OC use and by 20 years is almost 50% below that in women who have never used OCs (37).

When taken for more than 12 months, combined OCs confer equal protection against the three major histological subtypes of endometrial cancer: adenocarcinoma, adenosquamous carcinoma, and adenoacanthoma (38). Whereas earlier studies evaluated the effect of high dose OC preparations, a Swedish case-control study indicates that lower dose (30–35 μ g) OC formulations provide comparable protection (39). DMPA has been shown to decrease the risk of endometrial cancer by 80%, with protective effects lasting for at least eight years after cessation of treatment (40).

Ovarian Cancer

The risk of ovarian cancer declines with increasing duration of OC use. The incidence is 41%, 54%, and 61% lower with use for four years, eight years, and 12 years, respectively ($P < 0.0001$) (37). The protective effects of OCs become apparent after as little as three to six months of use and continue for up to 20 years after discontinuation (41, 42). The protective effect most likely derives from the progestin component of the combined OC (43). Risk for the four main histologic subtypes of epithelial ovarian cancer (serous, endometrioid, mucinous, and clear cell) is reduced to the same degree. Ovarian cancer risk associated with OCs containing the newer progestins, biphasic and triphasic pills, or lower dose OCs (20 μ g ethinyl estradiol) has not yet been clearly defined but appears similar.

Oral contraceptives may also have protective benefits for women at risk for hereditary ovarian cancer (44). Continuous use for 10 years in women with a family history of ovarian cancer may reduce the risk of epithelial ovarian cancer to a level less than or equal to that observed in women with no family history of the disease (45). A case-control study involving 207 women with hereditary ovarian cancer (with BRCA1 and BRCA2 genetic mutations) and 161 of their sisters as controls found significant protection against ovarian cancer with any past use of OCs compared with never use ($RR = 0.5$; 95% CI, 0.3–0.8) (44). Greatest protection was noted with six or more years of use ($RR = 0.3$; 95% CI, 0.1–0.7).

Cervical Cancer

The use of OCs has been associated with an increased risk of cervical intra-epithelial neoplasia (CIN) and cervical cancer (41, 46–49). However, since the human papillomavirus (HPV) has been implicated as the main causative agent in cervical cancer (50), OC use most likely acts as a co-factor in the development of this disease (51). A recent study of pooled data from case-control studies of invasive cervical cancer (eight studies) and carcinoma in situ (two studies) evaluated the risk of cervical cancer in women who were

using hormonal contraception and tested positive for the HPV (49). Hormonal contraception for fewer than five years did not increase the risk of cervical cancer ($OR = 0.73$; 95% CI, 0.52–1.03). However, risk increased with use of hormonal contraception for five to nine years ($OR = 2.82$; 95% CI, 1.46–5.42) and was greatest with use for 10 years or longer ($OR = 4.03$; 95% CI, 2.09–8.02). The proposed mechanism for this association is a hormonal effect on the cervix, a hypothesis that is supported by the increased risk of cervical cancer associated with increased parity and HPV (52) and the correlation between higher estradiol receptor concentrations induced by OCs and the risk of low-grade CIN (48).

Colorectal Cancer

There is growing epidemiologic evidence that OCs may protect women from developing colorectal cancer. In a meta-analysis, the overall estimated relative risk of colon cancer in OC users was 0.82 (95% CI, 0.74–0.92). The protection was stronger for women who had used OCs within the previous 10 years ($RR = 0.46$; 95% CI, 0.30–0.71) (53).

MEDICAL RISKS OF HORMONAL CONTRACEPTION

Hormonal Contraception and Bone Density

Depot-medroxyprogesterone acetate (DMPA) use has been associated with short-term bone loss in reproductive age women, an observation attributed to lower ovarian estrogen production resulting from suppression of gonadotropin secretion. Cross-sectional studies have demonstrated a decrease in bone density in the spine and distal radius in women currently using DMPA (54–56). Although bone loss correlates with duration of treatment and becomes significant after 15 years of use (57), other studies indicate the effect is reversible after discontinuation of treatment and therefore may not pose any longer term risk (55, 58, 59). Bone loss has also been documented in a small study of adolescent girls using DMPA; a 3% decrease in lumbar spine bone mineral was observed in users compared to a 9% increase in controls (60). These observations have raised legitimate concerns that DMPA use in adolescents may prevent them from achieving normal peak bone mass, although the contraceptive benefits of DMPA use in the adolescent population may outweigh any potential adverse effect on bone density.

The levonorgestrel-containing contraceptive implant has also been evaluated for its effect on bone density, although data are limited. A small study in seven adolescents showed a significant increase in lumbar spine bone density over 12 months (60). Studies of forearm bone density in adults have yielded conflicting results, with both increases and decreases reported (55, 61). The levonorgestrel implant should not decrease bone density as much as DMPA, because estrogen levels are not as consistently suppressed (18), and 19-norprogesterogens have a beneficial effect on bone (62).

The effect of combination OCs on bone density is less clear. Studies in postmenopausal women who have previously used OCs have revealed improved bone density that correlates with years of use (63). Reduced fracture risks that correlate with the use of high-dose formulations and use after the age of 40 have also been demonstrated in postmenopausal women (64). Short-term studies in current users have demonstrated a small gain in bone density or no significant change (55, 56, 65). One small study suggested that gain in bone density over a 12-month period was greater with a norethindrone-containing OC compared to another containing desogestrel (56). However, no study has shown a detrimental effect of OCs on bone mass.

Myocardial Infarction

Myocardial infarction is extremely rare among reproductive aged women. The base line risk of myocardial infarction among healthy women rises from two per million at age 30 to 34 to 20 per million at age 40 to 44 (66). Use of low dose OCs increases the risk of myocardial infarction by approximately two-fold among users even after controlling for cardiovascular risk factors (including smoking, hypertension, hypercholesterolemia, diabetes, and obesity) (67). The myocardial infarction risk in OC users is increased by smoking, an effect that is more noticeable among women over age 35 years. For women under age 35, the incidence of myocardial infarction for smokers (35 per million) is increased 10-fold over that for nonsmokers (3 per million). For women over age 35, the risks of myocardial infarction are significantly higher for both smoking (40 per 100,000) and non-smoking women (3 per 100,000) (68). Oral contraceptives should therefore be used with caution in women over age 35 who smoke.

Ischemic Stroke

Ischemic stroke too is very rare among healthy reproductive aged women. The annual incidence rises with increasing age (6 per million at age 20–24, 10 per million at age 30–34, and 16 per million at age 40–44) (66). A summary of five epidemiological, case–control studies involving 257 exposed cases estimated that the risk of ischemic stroke was 2.2-fold higher (95% CI, 1.9–2.7) with current use of OCs containing <50 µg ethinyl estradiol, compared to non-users. Risk is not related to the progestin component of OCs (69). Among women with a history of migraines, the relative risk of ischemic stroke is not significantly increased for those using OCs containing <50 µg ethinyl estradiol, compared to non-users (70, 71). The risk of ischemic stroke associated with OCs is not significantly different in women with simple migraine (no aura) compared to those with classic migraine (with aura) (70, 71). The risk of stroke in women with migraines who use OCs is increased significantly by smoking (70). Oral contraceptives should not be used in patients with visual changes or focal neurologic deficits associated with migraine headache (i.e., weakness, speech deficits).

Hemorrhagic Stroke

The incidence of hemorrhagic stroke among healthy women rises from 13 to 24 to 46 cases per million per year for ages 20 to 24, 30 to 34, and 40 to 44 years, respectively (66). In a review of three studies completed in the 1990s, the OC-associated risk of hemorrhagic stroke for users under 35 years of age was not increased compared to non-users (RR = 1.0, 95% CI, 0.7–1.5). For OC users ages 35 and older, the OC-associated risk of hemorrhagic stroke was 2.2-fold higher compared with non-users (95% CI, 1.5–3.3).

Venous Thromboembolism

The incidence of venous thromboembolism (VTE) among healthy women is low but increases with age from 32 to 46 to 59 cases per million per year for ages 20 to 24, 30 to 34, and 40 to 44 years, respectively (66). Oral contraceptive use is associated with a three-fold higher risk of VTE (66). The risk appears to be proportional to the estrogen dose (72). However, studies have also suggested that the risk of VTE is less than two-fold higher for OC formulations containing the newer progestins (desogestrel and gestodene compared to levonorgestrel (73). A meta-analysis including three cohort and nine case–control studies of VTE risk associated with newer progestins yielded an overall adjusted odds ratio (OR) for VTE of 1.7 (95% CI, 1.4–2.0) with an absolute excess risk of 1.5 events per 10,000 woman-years for women using formulations containing desogestrel or gestodene (74). The risk remained significantly elevated for short-term and long-term users and for users of differing ages but is far less than that associated with pregnancy. Although the risk of VTE is statistically significant, it is still a rare event and would translate to one to two cases per 10,000 women years of use (75). The FDA has suggested no change in prescribing (57).

The risk of VTE associated with prothrombotic conditions is further increased with OC use. The most common of these is the Factor V Leiden mutation which results in resistance to activated protein C. The risk of VTE in women who are heterozygous for the mutation is seven-fold higher than in non-carriers. In women who are heterozygous carriers and use OCs, the risk is 35-fold higher (95% CI, 7.8–154) (76). The prevalence of the Factor V Leiden mutation is 5% in Caucasians but extremely low in Asian and African populations (77). A genetic defect in prothrombin is also associated with increased risk of VTE and increases the risk of VTE in users of OCs (78). Other known factors involved in the development of VTE include protein C, protein S, and antithrombin III deficiencies (76). The presence of more than one of these mutations increases the risk of VTE significantly. Oral contraceptives should not be used in individuals who carry these mutations. At this time, screening for these mutations prior to initiating OC use is recommended only for women having a personal or family history of thrombosis (76).

CONCLUSIONS

- Hormonal contraception is a safe and effective form of reversible contraception.
- New transdermal, injectable, vaginal, and implantable delivery methods appear to have safety and efficacy comparable to that of OCs.
- Modifications of OCs, including progressively lower doses of estrogens and the development of newer progestins, have maintained contraceptive efficacy while reducing the incidence of side effects.
- Emergency contraception using either combined estrogen/progestin or progestin-only regimens is effective in reducing the risk of pregnancy. The progestin-only regimen may be more effective with fewer side effects.
- There is a reduced incidence of endometrial and ovarian cancers in past and current users of OCs.
- There is a potential for increased risk of cervical cancer with longer durations of OC use.
- There is no increase in lifetime risk of breast cancer associated with OC use, although prevalence is modestly increased in the first five years of use.
- The risk of stroke and myocardial infarction in reproductive aged women is low, but is increased in OC users over age 35 who smoke.
- The risk of VTE is increased in OC users and higher in those heterozygous for mutations resulting in prothrombotic conditions. At this time, screening for thrombophilias prior to initiating therapy with OCs is recommended only for women having a personal or a family history of thrombosis.

Acknowledgments: This report was developed under the direction of the Practice Committee of the American Society for Reproductive Medicine as a service to its members and other practicing clinicians. While this document reflects appropriate management of a problem encountered in the practice of reproductive medicine, it is not intended to be the only approved standard of practice or to dictate an exclusive course of treatment. Other plans of management may be appropriate, taking into account the needs of the individual patient, available resources, and institutional or clinical practice limitations. The Practice Committee and the Board of Directors of the American Society for Reproductive Medicine approved this report in February 2004.

References

1. Speroff L, Glass RH, Kase NG, eds. Clinical gynecologic endocrinology and infertility. Baltimore: 1999;867-945.
2. Sang GW, Shao QX, Ge RS, Ge JL, Chen JK, Song S, et al. A multicenter phase III comparative clinical trial of MesiGyna, Cyclofem and Injectable No. 1 given by intramuscular injection to Chinese women. II. The comparison of bleeding patterns. Contraception 1995; 51:185-92.
3. Kaunitz AM, Garceau RJ, Cromie MA. Comparative safety, efficacy, and cycle control of Lunelle monthly contraceptive injection (medroxyprogesterone acetate and estradiol cypionate injectable suspension) and Ortho-Novum 7/7/7 oral contraceptive (norethindrone/ethinyl estradiol triphasic). Lunelle Study Group. Contraception 1999;60:179-87.
4. Garceau RJ, Wajszczuk CJ, Kaunitz AM, Lunelle Study Group. Bleeding patterns of women using Lunelle™ monthly contraceptive injections (medroxyprogesterone acetate and estradiol cypionate injectable suspension) compared with those of women using Ortho-Novum® 7/7/7 (norethindrone/ethinyl estradiol triphasic) or other oral contraceptives. Contraception 2000;62:289-95.
5. Rahimy MH, Ryan KK. Lunelle monthly contraceptive injection (medroxyprogesterone acetate and estradiol cypionate injectable suspension): assessment of return of ovulation after three monthly injections in surgically sterile women. Contraception 1999;60:189-200.
6. Abrams LS, Skee D, Natarajan J, Wong FA. Pharmacokinetic overview of Ortho Evra/Evra. Fertil Steril 2002;77(Suppl 2):S3-12.
7. Ziemann M, Guillebaud J, Weisberg E, Shangold GA, Fisher AC, Creasy GW. Contraceptive efficacy and cycle control with the Ortho Evra/Evra transdermal system: the analysis of pooled data. Fertil Steril 2002; 77(Suppl 2):S13-8.
8. Audet MC, Moreau M, Koltun WD, Waldbaum AS, Shangold G, Fisher AC, ORTHO EVRA/EVRA 004 Study Group. Evaluation of contraceptive efficacy and cycle control of a transdermal contraceptive patch vs an oral contraceptive: a randomized controlled trial. JAMA 2001; 285:2347-54.
9. Mulders TM, Dieben TO. Use of novel combined contraceptive vaginal ring NuvaRing for ovulation inhibition. Fertil Steril 2001;75:865-70.
10. Roumen FJ, Apter D, Mulders TM, Dieben TO. Efficacy, tolerability and acceptability of a novel contraceptive vaginal ring releasing etonogestrel and ethinyl estradiol. Hum Reprod 2001;16:469-75.
11. Luukkainen T, Pakarinen P, Toivonen J. Progestin-releasing intrauterine systems. Semin Reprod Med 2001;19:355-63.
12. Andersson K, Odland V, Rybo G. Levonorgestrel-releasing and copper-releasing (Nova T) IUDs during five years of use: a randomized comparative trial. Contraception 1994;49:56-72.
13. Nilsson CG. Comparative quantitation of menstrual blood loss with a d-norgestrel-releasing IUD and a Nova-T-copper device. Contraception 1977;15:379-87.
14. Hidalgo M, Bahamondes L, Perrotti M, Diaz J, Dantas-Monteiro C, Petta C. Bleeding patterns and clinical performance of the levonorgestrel-releasing intrauterine system (Mirena) up to two years. Contraception 2002;65:129-32.
15. Sivin I, Stern J, Diaz S, Pavez M, Alvarez F, Brache V, et al. Rates and outcomes of planned pregnancy after use of Norplant capsules, Norplant II rods, or levonorgestrel-releasing or copper TCu 380Ag intrauterine contraceptive devices. Am J Obstet Gynecol 1992;166:1208-13.
16. Sivin I, Campodonico I, Kiriwat O, Holma P, Diaz S, Wan L, et al. The performance of levonorgestrel rod and Norplant contraceptive implants: a 5 year randomized study. Hum Reprod 1998;13:3371-8.
17. Sivin I, Alvarez F, Mishell DR Jr, Darney P, Wan L, Brache V, et al. Contraception with two levonorgestrel rod implants. A 5-year study in the United States and Dominican Republic. Contraception 1998;58: 275-82.
18. Meckstroth KR, Darney PD. Implant contraception. Semin Reprod Med 2001;19:339-54.
19. Hammond GL, Rabe T, Wagner JD. Preclinical profiles of progestins used in formulations of oral contraceptives and hormone replacement therapy. Am J Obstet Gynecol 2001;185:S24-31.
20. Krattenmacher R. Drospirenone: pharmacology and pharmacokinetics of a unique progestogen. Contraception 2000;62:29-38.
21. Poindexter A. The emerging use of the 20-microg oral contraceptive. Fertil Steril 2001;75:457-65.
22. Rosenberg MJ, Meyers A, Roy V. Efficacy, cycle control, and side effects of low- and lower-dose oral contraceptives: a randomized trial of 20 micrograms and 35 micrograms estrogen preparations. Contraception 1999;60:321-9.
23. Miller L, Notter KM. Menstrual reduction with extended use of combination oral contraceptive pills: randomized controlled trial. Obstet Gynecol 2001;98:771-8.
24. Kwiecien M, Edelman A, Nichols MD, Jensen JT. Bleeding patterns and patient acceptability of standard or continuous dosing regimens of a low-dose oral contraceptive: a randomized trial. Contraception 2003; 67:9-13.
25. Piaggio G, von Hertzen H, Grimes DA, Van Look PF. Timing of emergency contraception with levonorgestrel or the Yuzpe regimen. Task Force on Postovulatory Methods of Fertility Regulation. Lancet 1999;353:721.
26. Trussell J, Rodriguez G, Ellertson C. New estimates of the effectiveness of the Yuzpe regimen of emergency contraception. Contraception 1998; 57:363-9.
27. Trussell J, Rodriguez G, Ellertson C. Updated estimates of the effectiveness of the Yuzpe regimen of emergency contraception. Contraception 1999;59:147-51.
28. Task Force on Postovulatory Methods of Fertility Regulation. Randomised controlled trial of levonorgestrel versus the Yuzpe regimen of combined oral contraceptives for emergency contraception. Lancet 1998;352:428-33.
29. Raymond EG, Creinin MD, Barnhart KT, Lovvorn AE, Rountree RW, Trussell J. Meclizine for prevention of nausea associated with use of emergency contraceptive pills: a randomized trial. Obstet Gynecol 2000;95:271-7.

30. Nielsen CL, Miller L. Ectopic gestation following emergency contraceptive pill administration. *Contraception* 2000;62:275–6.
31. American College of Obstetricians and Gynecologists. ACOG practice bulletin no. 25: emergency oral contraception. Washington, DC: American College of Obstetricians and Gynecologists, 2001.
32. Comparison of three single doses of mifepristone as emergency contraception: a randomised trial. Task Force on Postovulatory Methods of Fertility Regulation. *Lancet* 1999;353:697–702.
33. Ashok PW, Stalder C, Wagaarachchi PT, Flett GM, Melvin L, Templeton A. A randomised study comparing a low dose of mifepristone and the Yuzpe regimen for emergency contraception. *BJOG* 2002;109:553–60.
34. Glasier A, Thong KJ, Dewar M, Mackie M, Baird DT. Mifepristone (RU 486) compared with high-dose estrogen and progestogen for emergency postcoital contraception. *N Engl J Med* 1992;327:1041–4.
35. Ries LAG, Eisner MP, Kosary CL, Hankey BF, Miller BA, Clegg L, et al. SEER cancer statistics review, 1975–2000. Bethesda, MD: National Cancer Institute, 2003.
36. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and hormonal contraceptives: collaborative reanalysis of individual data on 53,297 women with breast cancer and 100,239 women without breast cancer from 54 epidemiological studies. *Lancet* 1996;347:1713–27.
37. Schlesselman JJ, Collins JA. The influence of steroids on gynecologic cancers. In: Fraser IS, Jansen RPS, Lobo RA, Whitehead MI, eds. *Estrogens and progestogens in clinical practice*. London: Churchill Livingstone, 1999:831–64.
38. The Cancer and Steroid Hormone Study of the Centers for Disease Control and the National Institute of Child Health and Human Development. Combination oral contraceptive use and the risk of endometrial cancer. *JAMA* 1987;257:796–800.
39. Weiderpass E, Adami HO, Baron JA, Magnusson C, Lindgren A, Persson I. Use of oral contraceptives and endometrial cancer risk (Sweden). *Cancer Causes Control* 1999;10:277–84.
40. The WHO Collaborative Study of Neoplasia and Steroid Contraceptives. Depot-medroxyprogesterone acetate (DMPA) and risk of endometrial cancer. *Int J Cancer* 1991;49:186–90.
41. Schlesselman JJ. Net effect of oral contraceptive use on the risk of cancer in women in the United States. *Obstet Gynecol* 1995;85:793–801.
42. The Cancer and Steroid Hormone Study of the Centers for Disease Control and the National Institute of Child Health and Human Development. The reduction in risk of ovarian cancer associated with oral contraceptive use. *N Engl J Med* 1987;316:650–5.
43. Schildkraut JM, Calingaert B, Marchbanks PA, Moorman PG, Rodriguez GC. Impact of progestin and estrogen potency in oral contraceptives on ovarian cancer risk. *J Natl Cancer Inst* 2002;94:32–8.
44. Narod SA, Risch H, Moslehi R, Dorum A, Neuhausen S, Olsson H, et al. Oral contraceptives and the risk of hereditary ovarian cancer. Hereditary Ovarian Cancer Clinical Study Group. *N Engl J Med* 1998;339:424–8.
45. Gross TP, Schlesselman JJ. The estimated effect of oral contraceptive use on the cumulative risk of epithelial ovarian cancer. *Obstet Gynecol* 1994;83:419–24.
46. Ylitalo N, Sorensen P, Josefsson A, Frisch M, Sørensen P, Ponten J, et al. Smoking and oral contraceptives as risk factors for cervical carcinoma in situ. *Int J Cancer* 1999;81:357–65.
47. Kjellberg L, Hallmans G, Ahren AM, Johansson R, Bergman F, Wadell G, et al. Smoking, diet, pregnancy and oral contraceptive use as risk factors for cervical intra-epithelial neoplasia in relation to human papillomavirus infection. *Br J Cancer* 2000;82:1332–8.
48. Salazar EL, Sojo-Aranda I, Lopez R, Salcedo M. The evidence for an etiological relationship between oral contraceptive use and dysplastic change in cervical tissue. *Gynecol Endocrinol* 2001;15:23–8.
49. Moreno V, Bosch FX, Munoz N, Meijer CJ, Shah KV, Walboomers JM, et al. Effect of oral contraceptives on risk of cervical cancer in women with human papillomavirus infection: the IARC multicentric case-control study. *Lancet* 2002;359:1085–92.
50. Walboomers JM, Jacobs MV, Manos MM, Bosch FX, Kummer JA, Shah KV, et al. Human papillomavirus is a necessary cause of invasive cervical cancer worldwide. *J Pathol* 1999;189:12–9.
51. Skegg DC. Oral contraceptives, parity, and cervical cancer. *Lancet* 2002;359:1080–1.
52. Munoz N, Franceschi S, Bosetti C, Moreno V, Herrero R, Smith JS, et al. Role of parity and human papillomavirus in cervical cancer: the IARC multicentric case-control study. *Lancet* 2002;359:1093–101.
53. Fernandez E, La Vecchia C, Balducci A, Chatenoud L, Franceschi S, Negri E. Oral contraceptives and colorectal cancer risk: a meta-analysis. *Br J Cancer* 2001;84:722–7.
54. Cundy T, Cornish J, Roberts H, Elder H, Reid IR. Spinal bone density in women using depot medroxyprogesterone contraception. *Obstet Gynecol* 1998;92:569–73.
55. Petitti DB, Piaggio G, Mehta S, Cravioto MC, Meirik O. Steroid hormone contraception and bone mineral density: a cross-sectional study in an international population. The WHO Study of Hormonal Contraception and Bone Health. *Obstet Gynecol* 2000;95:736–44.
56. Berenson AB, Radecki CM, Grady JJ, Rickert VI, Thomas A. A prospective, controlled study of the effects of hormonal contraception on bone mineral density. *Obstet Gynecol* 2001;98:576–82.
57. American College of Obstetricians and Gynecologists. ACOG committee revises recommendations on OCs with third-generation progestins. *ACOG Today* 1998;July.
58. Cundy T, Cornish J, Evans MC, Roberts H, Reid IR. Recovery of bone density in women who stop using medroxyprogesterone acetate. *BMJ* 1994;308:247–8.
59. Orr-Walker BJ, Evans MC, Ames RW, Clearwater JM, Cundy T, Reid IR. The effect of past use of the injectable contraceptive depot medroxyprogesterone acetate on bone mineral density in normal postmenopausal women. *Clin Endocrinol* 1998;49:615–8.
60. Cromer BA, Blair JM, Mahan JD, Zibners L, Naumovski Z. A prospective comparison of bone density in adolescent girls receiving depot medroxyprogesterone acetate (Depo-Provera), levonorgestrel (Norplant), or oral contraceptives. *J Pediatr* 1996;129:671–6.
61. Naessen T, Olsson SE, Gudmundson J. Differential effects on bone density of progestogen-only methods for contraception in premenopausal women. *Contraception* 1995;52:35–9.
62. Horowitz M, Wishart JM, Need AG, Morris HA, Nordin BE. Effects of norethisterone on bone related biochemical variables and forearm bone mineral in post-menopausal osteoporosis. *Clin Endocrinol* 1993;39:649–55.
63. Kleerekoper M, Brienza RS, Schultz LR, Johnson CC. Oral contraceptive use may protect against low bone mass. Henry Ford Hospital Osteoporosis Cooperative Research Group. *Arch Intern Med* 1991;151:1971–6.
64. Michaelsson K, Baron JA, Farahmand BY, Persson I, Ljunghall S. Oral-contraceptive use and risk of hip fracture: a case-control study. *Lancet* 1999;353:1481–4.
65. Perotti M, Bahamondes L, Petta C, Castro S. Forearm bone density in long-term users of oral combined contraceptives and depot medroxyprogesterone acetate. *Fertil Steril* 2001;76:469–73.
66. Farley TM, Meirik O, Collins J. Cardiovascular disease and combined oral contraceptives: reviewing the evidence and balancing the risks. *Hum Reprod Update* 1999;5:721–35.
67. Tanis BC, van den Bosch MA, Kemmeren JM, Cats VM, Helmerhorst FM, Algra A, et al. Oral contraceptives and the risk of myocardial infarction. *N Engl J Med* 2001;345:1787–93.
68. WHO Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. Acute myocardial infarction and combined oral contraceptives: results of an international multicentre case-control study. *Lancet* 1997;349:1202–9.
69. Kemmeren JM, Tanis BC, van den Bosch MA, Bollen EL, Helmerhorst FM, van der Graaf Y, et al. Risk of Arterial Thrombosis in Relation to Oral Contraceptives (RATIO) study: oral contraceptives and the risk of ischemic stroke. *Stroke* 2002;33:1202–8.
70. Chang CL, Donaghy M, Poulter N. Migraine and stroke in young women: case-control study. World Health Organization Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. *BMJ* 1999;318:13–8.
71. Curtis KM, Chrisman CE, Peterson HB, WHO Programme for Mapping Best Practices in Reproductive Health. Contraception for women in selected circumstances. *Obstet Gynecol* 2002;99:1100–12.
72. Gerstman BB, Piper JM, Tomita DK, Ferguson WJ, Stadel BV, Lundin FE. Oral contraceptive estrogen dose and the risk of deep venous thromboembolic disease. *Am J Epidemiol* 1991;133:32–7.
73. World Health Organization Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. Effect of different progestagens in low oestrogen oral contraceptives on venous thromboembolic disease. *Lancet* 1995;346:1582–8.
74. Kemmeren JM, Algra A, Grobbee DE. Third generation oral contraceptives and risk of venous thrombosis: meta-analysis. *BMJ* 2001;323:131–4.
75. Pymar HC, Creinin MD. The risks of oral contraceptive pills. *Semin Reprod Med* 2001;19:305–12.
76. Vandenbroucke JP, Koster T, Briet E, Reitsma PH, Bertina RM, Rosendaal FR. Increased risk of venous thrombosis in oral-contraceptive users who are carriers of factor V Leiden mutation. *Lancet* 1994;344:1453–7.
77. Vandenbroucke JP, Rosing J, Bloemenkamp KW, Middeldorp S, Helmerhorst FM, Bouma BN, et al. Oral contraceptives and the risk of venous thrombosis. *N Engl J Med* 2001;344:1527–35.
78. Martinelli I, Taioli E, Bucciarelli P, Akhavan S, Mannucci PM. Interaction between the G20210A mutation of the prothrombin gene and oral contraceptive use in deep vein thrombosis. *Arterioscler Thromb Vasc Biol* 1999;19:700–3.